

06-14-00

A

UTILITY PATENT APPLICATION TRANSMITTAL
(Small Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.
0152.00372

Total Pages in this Submission

TO THE ASSISTANT COMMISSIONER FOR PATENTS**Box Patent Application**
Washington, D.C. 20231

Transmitted herewith for filing under 35 U.S.C. 111(a) and 37 C.F.R. 1.53(b) is a new utility patent application for an invention entitled:

SERTOLI CELLS AS BIOCHAMBERS

and invented by:

Don F. Cameron; Paul R. Sanberg; Samuel Saporta; Joel J. HushenIf a **CONTINUATION APPLICATION**, check appropriate box and supply the requisite information:☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: _____

Which is a:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: _____

Which is a:

☒ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No.: _____

Enclosed are:

Application Elements

1. ☒ Filing fee as calculated and transmitted as described below
2. ☒ Specification having 28 pages and including the following:

- a. ☒ Descriptive Title of the Invention
- b. ☐ Cross References to Related Applications (if applicable)
- c. ☒ Statement Regarding Federally-sponsored Research/Development (if applicable)
- d. ☐ Reference to Microfiche Appendix (if applicable)
- e. ☒ Background of the Invention
- f. ☒ Brief Summary of the Invention
- g. ☒ Brief Description of the Drawings (if drawings filed)
- h. ☒ Detailed Description
- i. ☒ Claim(s) as Classified Below
- j. ☒ Abstract of the Disclosure

CERTIFICATE OF MAILING BY "EXPRESS MAIL"

"EXPRESS MAIL" Mailing Label Number **EL 405 594 443 US**Date of Deposit **6-13-00**

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office To Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.


(Signature of person mailing paper or fee)

**UTILITY PATENT APPLICATION TRANSMITTAL
(Small Entity)**

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.
0152.00372

Total Pages in this Submission

Application Elements (Continued)

3. ☒ Drawing(s) *(when necessary as prescribed by 35 USC 113)*

a. ☐ Formal b. ☒ Informal Number of Sheets 6

4. ☒ Oath or Declaration

a. ☐ Newly executed *(original or copy)* ☒ Unexecuted

b. ☐ Copy from a prior application (37 CFR 1.63(d)) *(for continuation/divisional application only)*

c. ☒ With Power of Attorney ☐ Without Power of Attorney

d. ☐ DELETION OF INVENTOR(S)

Signed statement attached deleting inventor(s) named in the prior application,
see 37 C.F.R. 1.63(d)(2) and 1.33(b).

5. ☐ Incorporation By Reference *(usable if Box 4b is checked)*

The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered as being part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

6. ☐ Computer Program in Microfiche

7. ☐ Genetic Sequence Submission *(if applicable, all must be included)*

a. ☐ Paper Copy

b. ☐ Computer Readable Copy

c. ☐ Statement Verifying Identical Paper and Computer Readable Copy

Accompanying Application Parts

8. ☐ Assignment Papers *(cover sheet & documents)*

9. ☐ 37 CFR 3.73(b) Statement *(when there is an assignee)*

10. ☐ English Translation Document *(if applicable)*

11. ☐ Information Disclosure Statement/PTO-1449 ☐ Copies of IDS Citations

12. ☐ Preliminary Amendment

13. ☒ Acknowledgment postcard

14. ☒ Certificate of Mailing

☐ First Class ☒ Express Mail *(Specify Label No.):* EL 405 594 443 US

06/13/00



USPTO

UTILITY PATENT APPLICATION TRANSMITTAL (Small Entity)

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Docket No.
0152.00372

Total Pages in this Submission

Accompanying Application Parts (Continued)

15. ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
16. ☒ Small Entity Statement(s) - Specify Number of Statements Submitted: 1

17. ☐ Additional Enclosures (please identify below):


Fee Calculation and Transmittal

CLAIMS AS FILED

For	#Filed	#Allowed	#Extra	Rate	Fee
Total Claims	24	- 20 =	4	x \$9.00	\$36.00
Indep. Claims	7	- 3 =	4	x \$39.00	\$156.00
Multiple Dependent Claims (check if applicable) <input type="checkbox"/>					\$0.00
BASIC FEE					\$345.00
OTHER FEE (specify purpose)					\$0.00
TOTAL FILING FEE					\$537.00

- ☒ A check in the amount of \$537.00 to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. 11-1449 as described below. A duplicate copy of this sheet is enclosed.
- ☐ Charge the amount of as filing fee.
- ☒ Credit any overpayment.
- ☒ Charge any additional filing fees required under 37 C.F.R. 1.16 and 1.17.
- ☐ Charge the issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

Dated: June 13, 2000


Signature
Amy E. Rinaldo, Reg. No. 45,791
KOHN & ASSOCIATES
30500 Northwestern Highway, Suite 410
Farmington Hills, Michigan 48334
(248) 539-5050
FAX: (248) 539-5055

cc:

PATENT

Attorney's Docket Number: 0152.00372

Applicant or Patentee: Cameron et al.
Serial or Patent No: _____
Filed or Issued: _____
Title: SERTOLI CELLS AS BIOCHAMBERS

VERIFIED STATEMENT CLAIMING SMALL ENTITY STATUS
[37 CFR 1.9(f) and 1.27(c)]-SMALL BUSINESS CONCERN

I hereby declare that I am:

- _____ the owner of the small business concern identified below:
☒ an official of the small business concern empowered to
act on behalf of the concern identified below:

Name of Concern: University of South Florida Research Foundation, Inc.
Address of Concern: 4202 East Fowler Avenue - FAO 126
Tampa, Florida 33620-4962

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 13 CFR 121.3-18, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees under Section 41(a) and (b) of Title 35, United States Code, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement: (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when, either directly or indirectly, one concern controls or has the power to control the other, or a third-party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention described in:

- ☒ the specification filed herewith with title as listed above.
_____ the application identified above.
_____ the patent identified above.

If the rights held by the above-identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below* and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c), if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

* NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

Each such person, concern or organization having any rights in the invention is listed below:

_____ No such person, concern, or organization exists.

 X Each such person, concern or organization is listed below.

NAME: University of South Florida

ADDRESS: 4202 East Fowler Avenue - FAO 126

Tampa, Florida 33620-4962

 Individual Small Business X Nonprofit Organization

NAME: _____

ADDRESS: _____

 Individual Small Business Nonprofit Organization

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. [37 CFR 1.28(b)]

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Name of Person Signing: Kenneth G. Preston

Title in Organization: Executive Director

Address of Person Signing: 4202 East Fowler Avenue - FAC 126

Tampa, Florida 33620-4962

SIGNATURE:  Date: 6/1/00

115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933

* NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27)

NAME: University of South Florida Research Foundation, Inc.

ADDRESS: 4202 East Fowler Avenue - FAO 126

Tampa, Florida 33620-4962

 Individual X Small Business Nonprofit Organization

NAME : _____

ADDRESS: _____

Individual Small Business Nonprofit Organization

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. [37 CFR 1.28(b)]

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

Name of Person Signing: George R. Newkome, Ph.D.

Title in Organization: Vice President for Research

Address of Person Signing: 4202 East Fowler Avenue - FAO 126

Tampa, Florida 33620-4962

SIGNATURE: Date. 6/1/00

SERTOLI CELLS AS BIOCHAMBERS

GRANT INFORMATION

5

Supported in part by NASA Grant NAG8-1381.

BACKGROUND OF THE INVENTION

1. TECHNICAL FIELD

10

The present invention relates to methods of transplanting cells. More specifically, the present invention relates to methods of transplanting cells to create a localized immunosuppressive effect in the tissue receiving the transplanted cells.

15

2. BACKGROUND ART

20

The central nervous system (CNS) has poor regenerative capacity which is exemplified in a number of neurodegenerative disorders. An example of such a disorder is Parkinson's disease. The preferred pharmacotherapy for Parkinson's disease is the administration of L-dopa which slows the progression of this disease in some humans. However, the neuropathological damage and the consequent behavioral deficits is not reversed by this treatment protocol.

25

30

Laboratory and clinical studies have shown that the transplantation of cells into the CNS is a potentially significant alternative therapeutic modality for neurodegenerative disorders such as Parkinson's disease (Victorin et al., 1990; Lindvall et al., 1990; Sanberg et al., 1994; Bjorlund and Stenevi, 1985; Freeman et al., 1994). In some cases, transplanted neural tissue can survive

and form connections with the CNS of the recipient (i.e. the host). When successfully accepted by the host, the transplanted tissue (i.e. the graft) has been shown to ameliorate the behavioral deficits associated with the disorder (Victorin et al., 1990). The obligatory step for the success of this kind of treatment is the prevention of graft rejection (i.e. graft acceptance).

Currently, fetal neural tissue is the primary graft source for neural transplantation (Lindvall et al., 1990; Bjorklund, 1992; Isacson et al., 1986; Sanberg et al., 1994). Other viable graft sources include adrenal chromaffin cells and various cell types that secrete nerve growth factors and trophic factors. The field of neural tissue transplantation as a productive treatment protocol for neurodegenerative disorders has received much attention resulting in its progression to clinical trials. Preliminary results and clinical observations are promising although the graft rejection phenomenon remains problematic.

Transplantation is also a valuable therapy for other diseases, such as insulin dependent diabetes mellitus. Insulin dependent diabetes mellitus is a major health problem. Current forms of therapy are not efficient and do not necessarily lead to a prevention of diabetic complications such as renal failure or blindness. A desirable treatment alternative is to provide the diabetic with an endogenous source of insulin, transplanting either the whole pancreas or the endocrine component of the pancreas (i.e. islets of Langerhans) into the diabetic recipient. Although, whole pancreas transplantation is successfully achieved with at least 60% of the grafts still functioning after transplantation for one year, a major weakness of this approach is the need for continuous immunosuppression with powerful and toxic immunosuppressant drugs.

5 The transplantation of the isolated islets containing the insulin secreting β -cells has received much attention in both animal models of diabetes (1-7) and in humans (8-16). However, islet transplantation to a variety of organ sites has met with little success as a viable treatment for diabetes. For example, islet transplantation of major histocompatibility complex (MHC) in the BB/W rat with spontaneous diabetes mellitus of autoimmune etiology results in destroyed islets within a few days by a recurrence of the autoimmune disease (17). Likewise, destruction of grafted cells in the diabetic BB/W rat occurs in grafted islets of MHC-incompatible donors (18, 19). In the course of finding a suitable organ or tissue site for islet transplantation, it was discovered that the relocated abdominal testis, in particular, provides an extraordinary safe environment for extended survival of islet grafts and some relief of the diabetic complications (20-22).

15 The testis has long been considered to be an immunologically privileged site (23-26) although the precise mechanism(s) by which it protects (suppresses) graft rejection has not been clearly defined. Isolated islets of MHC-compatible donors have been shown to survive for extended periods of time in the non-immunosuppressed BB/W rat if implanted in the rat's testis which is then placed into the host's abdominal cavity (20-22,27). Although the maintenance of functional islets allografts is significant, a more difficult task and far more potentially significant accomplishment, in terms of clinical applicability, is the induction of normoglycemia in diabetic animals by the implantation of cross-species islet xenografts.

25 Selawry and co-workers demonstrated the feasibility of such a procedure by successfully implanting incubated hamster islets into the BB/W rat abdominal testes (22,27,28). As a result of the abdominal testis/islet implant, the diabetic animals in these studies became normoglycemic. Long-

term survival of the islet xenografts did not require prolonged immunosuppression to prevent rejection and to maintain normal sugar levels. In all cases implant viability required the protective milieu of the abdominal testis. It now appears that the donor origin of these isolated islets does not
5 seem to influence their long-term survival. Islet cells grafted against major histocompatibility barriers (21), islet xenografts (27) and islets of MHC-compatible donors grafted into the testes of the diabetic BB/W rats functioned indefinitely in the recipient rendering the once diabetic animal normoglycemic.

10 The major weakness of this type of islet transplantation protocol is associated with the use of such an unconventional organ site. One major concern is the possibility of malignant transformation of germ cells at the higher core body temperature (29). More importantly, it would not be possible to use this transplantation protocol for the treatment of female diabetics.

15 Histological examination of grafted abdominal testes has shown that the islet implants are always found within the interstitial compartment of the gonad, which consists of the endocrine cells of Leydig, macrophages, blood vessels, testicular interstitial fluid and extracellular macromolecules (31). Any
20 of the secretory products of these cells are potentially capable of inhibiting the immune response. For instance, Born and Wekerle (32, 33) showed that active suppression of immune responses occurred by Leydig cells *in vitro*. These investigators speculated that the Leydig cells might prevent lymphocyte proliferative responses by creating an "immunologically neutral
25 zone" around the seminiferous tubules and thus decreasing the danger of T-cell infiltration in to the intratubular spaces. It was shown by Williams (34) that leukemic cells accumulate in the interstitial compartment where they are apparently protected against destruction by the host's immune defenses.

5 The "zone of protection" theory of Born and Wekerle (32) is attractive but it is not likely that this major component of the testicular interstitium, i.e. Leydig cells, is responsible for the synthesis of some protective (immunosuppressant) factor. Treatment of rats with ethane
10 dimethanesulphonate (EDS), which selectively destroys the Leydig cell completely, including steroidogenesis and all other functions, had no adverse effects on the survival of intratesticular islet allografts (30). It is not probable that germ cells were involved either, since these cells are readily depleted in the abdominal testis. By eliminating these cells, Cameron and Sewlary
15 concluded that the Sertoli cell was the most probable testicular cell type providing the testis with its unique immunologically privileged environment and that this cell was most likely responsible for the unexplained absence of islet rejection in abdominal testes (30). Based on these findings, Selawry and Cameron (35) attempted to create a similar immunologically privileged site outside of the testis utilizing Sertoli cells as an immunosuppressant agent. To
20 this end, isolated Sertoli cells were transplanted with isolated islets under the kidney capsule in female diabetic rats (see Figure 1). Results from this study showed this novel transplantation protocol resulted in normoglycemia and that long-term islet allograft survival was achieved in a traditionally immunologically hostile site. We concluded that the Sertoli cell, independent of the testicular milieu, secreted an immunosuppressant factor(s) which was neither androgenic nor inhibitory to ovulation since 6 of the 7 mated recipients became pregnant, carried a pregnancy to term and nursed the pups successfully (35).

25 For the long-term treatment of diabetes, it is clear that the presence of viable Sertoli cells is a prerequisite for long-term islet graft success and maintenance of long-term beta cell function. We do not yet clearly understand, however, the mechanism(s) which yield this observation. The
30 likely explanation is that the Sertoli cells secrete an immunosuppressant

factor(s) which cooperates with exogenous immunosuppressants such as cyclosporine A to prevent a complete immune response and subsequently tissue rejection (35). Sertoli cells are active secretory cell types synthesizing many proteins, some of which promote growth and others which have immunosuppressive capabilities (36, 55). Initial studies to verify such a factor have been positive to date. The effects of Sertoli cell conditioned media on Con A-stimulated spleen lymphocyte proliferation showed that products secreted by Sertoli cells inhibit lymphocyte proliferation in a dose-dependent manner. The synthesis was temperature dependent, occurring predominantly at 37°C and hormone dependent, requiring the presence of follicle stimulating hormone (FSH) in the Sertoli cell culture (see Figure 2). We further examined the mechanism of inhibition of lymphocyte proliferation and showed that preconditioned Sertoli cell media inhibited the production of the lymphokine IL-2 in a dose-dependent manner (see Figure 3A). Because the addition of exogenous IL-2 was not able to reverse this inhibition (see Figure 3B), it appears likely that the preconditioned media inhibited both IL-2 production and T-lymphocyte responsiveness to IL-2 (38) in concurrence with similar finding by DeCesarts et al. (39) It is widely acknowledged that all proliferating T-cells express IL-2 receptors, while resting cells do not, and that interaction of IL-2 with its receptor is an absolute requirement for the clonal expansion of activated T-cells (40). Because the prevention of IL-2 receptor interaction completely inhibits T-cell proliferation, we propose that both clonal expansion and viability of activated T-cells are suppressed by an immunosuppressive factor secreted by the Sertoli cells (35). In this fashion, the putative Sertoli cell derived immunosuppressant would appear to suppress the rejection by a mechanism similar to the action of cyclosporin A which also suppressed the production of IL-2 (41-44).

Although this hypothesis is appealing and with some research support of an indirect nature, it remains to be clearly unravelled. Recently, an additional and even more appealing hypothesis has received consideration attention. Bellgrau et al. (45*) in a letter to Nature showed that testis grafts that expressed Fas (CD95) ligand (FasL) survived indefinitely when transplanted under the kidney capsule, whereas testis grafts from *gld* mice (FasL deficient) were rejected when transplanted at the same site (45). A reverse transcriptase-polymerase chain reaction analysis demonstrated that Sertoli cells constitutively express FasL mRNA. Additionally, they showed that isolated Sertoli cells derived from normal, but not the *gld* mice survived indefinitely when transplanted under the kidney capsule. They concluded that the expression of functional FasL by Sertoli cells accounts for the immune-privilege nature of testis and suggested a mechanism by which Sertoli cells induce localized immune privilege to islets co-transplanted with Sertoli cells in an otherwise immune hostile site (i.e. subjacent the kidney capsule). They pointed out that FasL ligand-mediated immunosuppression would be expected to primarily target activated effector T cells rather than the activation steps that produce them, a mechanism by which Cyclosporin A produces immunosuppression. This would suggest that by targeting only activated T lymphocytes, grafted cell-associated FasL may provide a highly specific form of immunosuppression for ameliorating T-cell-dependent graft rejection. To this end, Lau et al, (46) transfected muscle cells with the FasL gene and co-transplanted them with islets beneath the kidney capsule and achieved local immunoprotection for the grafted islet, albeit for only 80 days. In a letter to Science, D. Green declared this a stunning advance and declared that "It's almost the Holy Grail of immunosuppression to restrict the suppression to the environment of the graft" (47). Selawry and Cameron (35) achieved the same results with long-term immunoprotection of the grafted islets and long-term maintenance of normoglycemia in the diabetic rat by co-transplanting the islets with the natural producer of FasL, Sertoli cells. The salient features of

terminally differentiated Sertoli cells that make them important and preferable as a transplantation facilitator are 1) they live for the life of the donor and may survive for the life of the recipient host (providing, thereby, long-term FAS-L induced local immunoprotection for the transplanted tissue or cells), 2) they do not divide and 3) they are easily isolated.

Since Sertoli cells secrete many growth enhancing factors including insulin-like growth factor I (55), the presence of Sertoli cells, in addition to their immunoprotective protective properties, may provide additional tropic and growth support to the transplant. Recently, Selawry et al, (48) showed that when cryopreserved pig Sertoli cells were thawed and immediately place in culture with Sertoli cells, there was a significant enhancement of post-thaw survival and insulin secretion when compared to thawed islets not co-cultured with Sertoli cells. They suggested that insulin-like growth factor I may have provided growth factor support to the cell membrane known to be damaged during freezing. Recently Sanberg et al (49-51) demonstrated that Sertoli cells can survive in the brain and, in fact, protect bovine adrenal chromaffin cell xenografts from rejection when co-transplanted into the striatum of the Parkinson's disease rat model. Even more significant, Sertoli cells alone transplanted into the PD rat result in the amelioration of motion dysfunction to the same degree as do chromaffin cells indicating a type of successful growth factor therapy, as yet unknown, provided for by the transplanted Sertoli cells (52). Similar to islet cells, Cameron et al (53) have shown that the post-thaw viability of fetal brain cells is significantly enhanced if the neuron are co-cultured with Sertoli cells again indicating the generalized ability of Sertoli cell secretory products to support the viability of isolated cells. For both islets and neurons, the growth and viability enhancing characteristics of Sertoli cells were evident only when the Sertoli cells were present as opposed to only media soluble factors found in expended pre-conditioned Sertoli cell media.

[illegible]

In general, systematic immunosuppression is necessary if successful transplantation is to be achieved in humans. Immunosuppression of the entire body (i.e. systemic) can result, eventually, in graft acceptance. It is acquired, however, by placing the individual at medical risk making the immunosuppressant therapy itself more of a liability than a benefit in some cases. For a lack of a better immunosuppressant treatment, systemic immunosuppressants, with Cyclosporine-A (CsA) as the treatment choice, have been used as adjunctive therapy in neural transplantation protocols (Sanberg et al., 1994; Freeman et al., 1994; Borlongan et al., 1995). Arguably, systemic CsA treatment may be contraproductive to successful graft acceptance in the CNS because of its systemic effect and because CsA itself has been shown to cause detrimental side effects and may in fact, be cytotoxic to neural tissues (Berden et al., 1985; deGroen et al., 1984).

It would be useful to develop a mechanism that enhances the productive cell transplantation techniques already utilized for neurodegenerative disorders, such as Parkinson's disease. This mechanism should improve these protocols in ways which would more effectively slow the neurodegenerative disease process, more actively promote the re-establishment of normal neural tissue physiology and better alleviate the functional disabilities associated with the neural tissue dysfunction. Likewise, it would be useful to provide trophic support for the transplanted cells. Further, it would be useful if this support lead to the reduction or elimination of systemic immunosuppression while maintaining the ability to immunosuppress locally (i.e. at the graft site) by an immunosuppressant which is biologically tolerated by the host. Sertoli cells may provide this desired option since it is clear from the diabetic studies, as summarized above, that co-transplantation with Sertoli cells will deliver local immunosuppression and promote, therefore, efficient graft acceptance and functional restoration of the tissue-related dysfunction.

SUMMARY OF THE INVENTION

According to the present invention, there is provided a biological chamber including outer walls of Sertoli cells and an inner lumen. Also provided is a transplantation facilitator including a biochamber which is formed from an engineered Sertoli tissue construct. A method of making biochambers by co-culturing facilitator cells and therapeutic cells is also provided. Additionally, there is provided a method of transplanting cells by incorporating therapeutic cells into a biochamber and transplanting the biochamber containing the therapeutic cells. Further, a method of treatment using these engineered biochambers is also included.

DESCRIPTION OF THE DRAWINGS

Other advantages of the present invention will be readily appreciated
5 as the same becomes better understood by reference to the following detailed
description when considered in connection with the accompanying drawings
wherein:

Figure 1 is a diagram showing the formation of a biochamber on a
10 substrate;

Figure 2 is a comparison showing the differences between the
conventional culture and a microgravity co-culture;

Figure 3 is a mechanism showing the way the Sertoli cells effect
15 immunosuppression at the graft site;

Figure 4 is a photograph showing Sertoli cells (SC) and islets (arrows)
in a Sertoli-islet tissue construct created in a conventional co-culture; B-cells
20 are immunostained for insulin;

Figure 5 is a photograph of Sertoli cells (SC) and B-cells (arrows) in a
Sertoli-islet tissue construct created in a conventional co-culture, B-cells are
immunostained for insulin;

Figure 6 is a photograph of Sertoli-Neuron-Aggregate-Cells (SNACS
for *in vitro* following coculture of rat Sertoli cells and NT2 neuros in simulated
microgravity utilizing the High Aspect Rotation Velocity (HARV) bioreactor;
and

Figure 7 is a photograph of Immunocytochemical staining of mouse FasL and human nuclear matrix proteins in Sertoli-Neuron Aggregated Cells (SNACs) following HARV incubated cocultures.

5

DETAILED DESCRIPTION OF THE INVENTION

Generally, the present invention provides a biological chamber system which is used for transplanting cells. More specifically, the biochamber is formed of facilitator cells such as, but not limited to, Sertoli cells which form a chamber or vessel having an inner cavity or lumen containing therein a population of cells different than the facilitator cells. In the preferred embodiment, this population of cells include therapeutic cells.

By "Biochamber" or "vessel", it is meant that a number of cells are engineered in such a manner as to form discrete walls about a lumen or center chamber. More specifically, the biochamber is formed by a structural modification of the Sertoli cells, this new structure being similar to the original Sertoli cell structure prior to cell harvesting. It is during this harvesting that the Sertoli cells are reorganized to form a central lumen in which the therapeutic cells are housed within a newly formed micro-environment. This micro-environment can contain therein therapeutic cells, which are used for transplantation. By "facilitator cell", as used herein, it is meant to include a cell which is able to provide localized immunosuppression or otherwise facilitate or make more effective the transplant. The facilitator cells provide bio-protection for the therapeutic or transplanted cells. This bio-protection includes, but is not limited to, protection from a biological source such as an immune response, whether cellular or humoral. In the preferred embodiment, the facilitator cell is a Sertoli cell. Such cells, as described hereinbelow, are able to reorganize to form walls defining an inner lumen. The biological/living

walls provide a physical as well as an immunological barrier for the cells contained therein. The apical secretions of Sertoli cells contribute to the unique trophic-bridge micro-environment of the luminal spaces in which therapeutic cells reside.

5

The term "therapeutic cell" as used herein, is meant to include the cells to be transplanted. For example, these therapeutic cells can include, but are not limited to, the following cells: dopaminergic cells, pancreatic islet cells, bovine chromaffin cells and immortalized neuron-like NT2 cells. The cells are therapeutic in that they can secrete hormones, factors, or the like that can have a therapeutic effect upon the host. They, like the Sertoli cell walls, are biosensitive in that they can respond to factors in their environment.

10

15

By modifying the harvested Sertoli cell by the methods of the instant invention that the cells reorganize into a tissue structure similar to that observed in the testis. They become a protective and nurturing barrier tissue, encapsulating the therapeutic cells in a unique micro-environment. Because the engineered Sertoli tissue construct captures the therapeutic cells in their new environment, this produces a dynamic support system for the therapeutic cells whereby the discreet units become efficient and viable within this special structure. Each biochamber becomes a discreet transplant unit, both nurtured and immunoprotected by the surrounding engineered Sertoli tissue.

20

25

In the preferred embodiment, Sertoli cells are isolated from a mammal, such as, but not limited to a prepubertal rat or pig testes and co-cultured with a therapeutic cell type in a culture environment that enhances tissue formation. This can be accomplished by co-culturing the different cell types in simulated microgravity culture utilizing the HARV bioreactor or other culture technologies. In a further embodiment, the co-culturing is performed without the microgravity environment.

30

002190-02926560

The addition of a basement membrane-like extracellular matrix to the incubation medium induces the epithelization and polarization of Sertoli cells, and subsequent formation of Sertoli-Sertoli junctional complexes between adjacent Sertoli cells, and the formation of a lumen or lumina. There is segregation of the Sertoli cells away from the therapeutic cells during the process of Sertoli cell epithelization leaving the therapeutic cells residing within the newly-formed luminal spaces. The luminal space(s) is/are created during this reorganization of the Sertoli cells and the formation of the Sertoli-Sertoli junctions. These junctions form an intraepithelial barrier similar to that observed in the testis and referred to as the blood-testis barrier. Apical polarization of Sertoli cell secretion is the likely mechanism by which the lumen is formed (Figures 1-3).. The reorganized Sertoli cells illustrated in Figures 1-3 create an item which is referred to as the Sertoli cell biochamber.

The Sertoli cell portion of the biochamber acts as a facilitator or a bridge cell for the transfer of material into and out of the lumen.

Examples of such biochambers, include but are not limited to, Islet-filled Sertoli cell biochambers (SICAs) and NT2 cell-filled biochambers (SNACs) which exemplify how therapeutic cell types can be incorporated into the Sertoli cell biochamber. SICAs secrete insulin in response to a glucose challenge (180 mg %) and also suppress activated lymphocytic proliferation (16). Similarly, SNACs enhance the differentiation of NT2 cells to the dopaminergic phenotype (17,18) and likewise provide for immunoprotection of the neurons as judged by the expression of FasL on the Sertoli cells (see Fig 3). SICAs and SNACs are therapeutic cell-filled Sertoli cell biochamber products created by this tissue engineering protocol and are designed for the use in therapeutic transplantation treatments for serious diseases such as diabetes and Parkinson's disease.

Since Sertoli cells are terminally differentiated, and the cells are mitotically inactive. They live for a long period of time, and potentially as long as any therapeutic cell type that can be engineered into the Sertoli cell biochamber. If transplanted in a Sertoli cell biochamber, therapeutic cells can be protected against immune surveillance and subsequent rejection in a micro-environment (provided for by Sertoli cell secreted growth and trophic factors) that also maintains and stimulates their functional phenotypes on a long-term basis. This has a significant impact on the successful transplantation treatment of many serious diseases and on the status of transplantation biology in general.

The above discussion provides a factual basis for the use of Sertoli cell biochambers. The methods used with and the utility of the present invention can be shown by the following non-limiting examples and accompanying figures.

EXAMPLES

GENERAL METHODS:

General methods in molecular biology: Standard molecular biology techniques known in the art and not specifically described were generally followed as in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York (1989), and in Ausubel et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Maryland (1989) and in Perbal, *A Practical Guide to Molecular Cloning*, John Wiley & Sons, New York (1988), and in Watson et al., *Recombinant DNA*, Scientific American Books, New York and in Birren et al (eds) *Genome Analysis: A Laboratory Manual Series, Vols. 1-4* Cold Spring Harbor Laboratory Press, New York (1998) and methodology as set forth in United

States patents 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057 and incorporated herein by reference. Polymerase chain reaction (PCR) was carried out generally as in *PCR Protocols: A Guide To Methods And Applications*, Academic Press, San Diego, CA (1990). In-situ (In-cell) PCR in combination with Flow Cytometry can be used for detection of cells containing specific DNA and mRNA sequences (Testoni et al, 1996, Blood 87:3822.)

General methods in immunology: Standard methods in immunology known in the art and not specifically described are generally followed as in Stites et al.(eds), *Basic and Clinical Immunology* (8th Edition), Appleton & Lange, Norwalk, CT (1994) and Mishell and Shiigi (eds), *Selected Methods in Cellular Immunology*, W.H. Freeman and Co., New York (1980).

Immunoassays

In general, immunocytochemistry ELISAs are the preferred immunoassays employed to assess a specimen. These assays are well known to those skilled in the art. Both polyclonal and monoclonal antibodies can be used in the assays. Where appropriate other immunoassays, such as radioimmunoassays (RIA) can be used as are known to those in the art.

Available immunoassays are extensively described in the patent and scientific literature. See, for example, United States patents 3,791,932; 3,839,153; 3,850,752; 3,850,578; 3,853,987; 3,867,517; 3,879,262; 3,901,654; 3,935,074; 3,984,533; 3,996,345; 4,034,074; 4,098,876; 4,879,219; 5,011,771 and 5,281,521 as well as Sambrook et al, *Molecular Cloning: A Laboratory Manual*, Cold Springs Harbor, New York, 1989

Delivery of gene products/therapeutics (compound):

The compound of the present invention is administered and dosed in accordance with good medical practice, taking into account the clinical

condition of the individual patient, the site and method of administration, scheduling of administration, patient age, sex, body weight and other factors known to medical practitioners. The pharmaceutically "effective amount" for purposes herein is thus determined by such considerations as are known in the art. The amount must be effective to achieve improvement including but not limited to improved survival rate or more rapid recovery, or improvement or elimination of symptoms and other indicators as are selected as appropriate measures by those skilled in the art.

The biochambers of the instant invention can be administered in various ways. These include subcutaneously or parentally, including intravenous, intraarterial, intramuscular, intraperitoneal and intranasal administration. Pharmaceutically acceptable carriers, diluents, adjuvants and vehicles are also useful for administration of the biochambers. These refer to any diluent, carrier, adjuvant or vehicle as commonly known to one of ordinary skill in the art.

EXAMPLE 1:

Recently, Sertoli cells have been utilized to facilitate islet transplantation on the basis that the testis-derived cells provide localized immunoprotection at the graft site and stimulate islet viability. The relationship between Sertoli cells and β -cells is not yet well defined *in vivo* nor *in vitro*. To further evaluated this relationship and to promote Sertoli/islet cell 3-dimensional aggregation (SICA) *in vitro*, Sertoli cells and islets were co-cultured in simulated microgravity using the NASA high aspect rotation velocity (HARV) bioreactor.

Sertoli cells, harvested from mammals by methods as known by those

of skill in the art, and islets, obtained by methods known to those of skill in the art as in Korbitt et al, were isolated from neonatal pigs by routine enzymatic digestion. Sertoli cells were placed immediately into HARVs at the time of isolation. Isolated islets were pre-cultured in flasks for 14 days (to expedite the removal of exocrine tissue) prior to incubation in HARV's with or without Sertoli cells. HARV co-cultures were incubated at 37° for 28 days in defined incubation medium consisting of DMEM; F-12 supplemented with ITS+ Retinol, and 1% Matrigel . Every 48 hours, 4 ml of media was removed and replaced with fresh media. The SICAs were exposed to a standard glucose challenge (180 mg% glucose) after which samples were collected every ten minutes for an hour and subsequently assayed for insulin by radioimmunoassay. Cell viability was determined by trypan blue exclusion, the presence of β -cells was determined by differential staining with dipherrylthiocarbazone and/or insulin immunostaining, and Sertoli cells were determined by FAS-L immunostaining.

By the end of the incubation period, Sertoli cells and islets had formed sizable (3-10 mm diameter) tissue constructs, with those formed in Matrigel mediums being larger. Cell viability was high (>80%) and β -cells were detected in both SICAs. In the three separate HARV incubations, the presence of Sertoli cells in SICA's enhanced the basal and total amount of insulin secreted in response to the glucose challenge when compared to islet-only HARV monocultures. In the presence of Sertoli cells, the SICA's insulin response to the elevated glucose was quicker and appeared to be prolonged.

EXAMPLE 2:

**FORMATION OF SERTOLI-NEURON AGGREGATED CELLS(SNACs) BY
SIMULATED MICROGRAVITY COCULTURE OF SERTOLI CELLS AND
IMORTALIZED NT2 CELLS**

[illegible][illegible]

	0	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99																																																																																																																																																												
0	00000000	00000001	00000010	00000011	00000100	00000101	00000110	00000111	00001000	00001001	00001010	00001011	00001100	00001101	00001110	00001111	00010000	00010001	00010010	00010011	00010100	00010101	00010110	00010111	00011000	00011001	00011010	00011011	00011100	00011101	00011110	00011111	00100000	00100001	00100010	00100011	00100100	00100101	00100110	00100111	00101000	00101001	00101010	00101011	00101100	00101101	00101110	00101111	00110000	00110001	00110010	00110011	00110100	00110101	00110110	00110111	00111000	00111001	00111010	00111011	00111100	00111101	00111110	00111111	01000000	01000001	01000010	01000011	01000100	01000101	01000110	01000111	01001000	01001001	01001010	01001011	01001100	01001101	01001110	01001111	01010000	01010001	01010010	01010011	01010100	01010101	01010110	01010111	01011000	01011001	01011010	01011011	01011100	01011101	01011110	01011111	01100000	01100001	01100010	01100011	01100100	01100101	01100110	01100111	01101000	01101001	01101010	01101011	01101100	01101101	01101110	01101111	01110000	01110001	01110010	01110011	01110100	01110101	01110110	01110111	01111000	01111001	01111010	01111011	01111100	01111101	01111110	01111111	10000000	10000001	10000010	10000011	10000100	10000101	10000110	10000111	10001000	10001001	10001010	10001011	10001100	10001101	10001110	10001111	10010000	10010001	10010010	10010011	10010100	10010101	10010110	10010111	10011000	10011001	10011010	10011011	10011100	10011101	10011110	10011111	10100000	10100001	10100010	10100011	10100100	10100101	10100110	10100111	10101000	10101001	10101010	10101011	10101100	10101101	10101110	10101111	10110000	10110001	10110010	10110011	10110100	10110101	10110110	10110111	10111000	10111001	10111010	10111011	10111100	10111101	10111110	10111111	11000000	11000001	11000010	11000011	11000100	11000101	11000110	11000111	11001000	11001001	11001010	11001011	11001100	11001101	11001110	11001111	11010000	11010001	11010010	11010011	11010100	11010101	11010110	11010111	11011000	11011001	11011010	11011011	11011100	11011101	11011110	11011111	11100000	11100001	11100010	11100011	11100100	11100101	11100110	11100111	11101000	11101001	11101010	11101011	11101100	11101101	11101110	11101111	11110000	11110001	11110010	11110011	11110100	11110101	11110110	11110111	11111000	11111001	11111010	11111011	11111100	11111101	11111110	11111111

NT2 cells. Some centrally located cells showed positive immunostaining for TH. It appeared that with MG, the Sertoli biochamber tissue construct was achieved with these two cell types, as described for the SICA (see Example 1). It is therefore concluded that the HARV coculture of Sertoli cells, and NT2 neurons with MG, resulted in the formation of NT2-filled Sertoli biochambers comprised of FasL positive Sertoli cells forming the biochamber wall and NuMa positive NT2 cells residing within the biochamber. The expression of TH suggests that some of the NT2 cells had differentiated into the dopaminergic phenotype indicating the use of these SNACs transplantation protocols for the treatment of experimental Parkinson's disease.

EXAMPLE 3

Isolated Sertoli cells from peripubertal rats and pancreatic islets from neonatal pigs were co-cultured by conventional culture technology in the same medium described for the HARV simulated microgravity coculture. Sertoli cells were pre-plated 48 hours on plastic or Matrigel substrates. Pre-treated isolated pig islets were added to the Sertoli cell-enriched monoculture 24 hours later. This Sertoli-Islet co-culture was incubated at 37°C and within 24 hr. islets had attached to and integrated into the underlying Sertoli cells. Within another 48-72hrs, Sertoli cells reorganized into spherical or chord-like aggregates. This process was enhanced for those co-cultures in which Sertoli cells had been plated on the Matrigel. Islets appeared to retain their structural integrity better in the non-Matrigel co-cultures (Figure 4) than in the cocultures not having a Matrigel substrate (Figure 5). Tissue constructs of Sertoli cells and pancreatic islet cells can be created in conventional coculture in a similar manner as that observed in simulated microgravity coculture.

Throughout this application, various publications, including United States patents, are referenced by author and year and patents by number. Full citations for the publications are listed below. The disclosures of these publications and patents in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art to which this invention pertains.

The invention has been described in an illustrative manner, and it is to be understood that the terminology which has been used is intended to be in the nature of words of description rather than of limitation.

Obviously, many modifications and variations of the present invention are possible in light of the above teachings. It is, therefore, to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described.

REFERENCES

1. Kaufman, et al. (1990).
- 5 2. Evans, et al. (1990).
3. Horaguchi, et al. (1981).
4. Kneteman, et al. (1990).
5. Ricordi, et al. (1990).
6. Thompson et al. (1990).
- 10 7. Calafiore, et al. (1990).
8. Gray, et al. (1984).
9. Scharp, et al. (1987).
10. Alderson, et al. (1987).
11. Ricordi, et al. (1989).
- 15 12. Scharp, D.W. (1988).
13. Kneteman, et al. (1986).
14. Warmock, et al. (1988).
15. Warnock, et al. (1989).
16. Kuhn, et al. (1985).
- 20 17. Naji, et al. (1981).
18. Weringer, et al. (1985).
19. Prowse, et al. (1986).
20. Selawry (1985).
- 25 21. Selawry et al. (1987).
22. Selawry, et al. (1986).
23. Whitmore, et al. (1978).
24. Head, et al. (1983).
25. Head, et al. (1983).

26. Hedger, M.P. (1989).
27. Barker et al. (1968).
28. Selawry, et al. (1989).
29. Martin, D.C. (1982).
- 5 30. Cameron, et al. (1990).
31. Fawcett, et al. (1973).
32. Born, et al. (1982).
33. Born, et al. (1981).
34. Williams, et al. (1978).
- 10 35. Selawry, et al. (1993).
36. Bardin, et al. (1988).
37. Griswold, M.D. (1993).
38. Selawry, et al. (1991).
39. DeCesarts, et al. (1992).
- 15 40. Cantrell, et al. (1984).
41. Leapman, et al. (1981).
42. Hess, A.D. (1985).
43. Green, et al. (1978).
44. Homan, et al. (1980).
- 20 45. Bellgrau, et al. (1995).
46. Lau, et al. (1996).
47. Wickelgren, L. (1996).
48. Selawry, et al. (1996).
49. Sanberg, et al. (1995).
- 25 50. Sanberg, et al. (1966).
51. Sanberg, et al. (1996).
52. Borlongan, et al. (1996).
53. Cameron, et al. (1996).
54. Gondos, et al. (1993).

55. Skinner, et al. (1993).
56. Edgington, S.M. (1992).
57. Goodwin, et al. (1993).
58. Goodwin, et al. (1993).
- 5 59. Goodwin, et al. (1992).
60. Becker et al. (1993).
67. Cameron et al. (1991).
68. Cameron, et al. (1993).
69. London, et al. (199).
- 10 70. Cameron et al. (1990).
71. Schwarz, et al. (1992).
72. Goodwin, et al. (1996).
73. Suda, et al. (1993).
74. Towbin, et al. (1979).
- 15 75. Prewett, et al. (1993).
76. Goodwin, et al. 91993).
77. Goodwin, et al. (1993).

20

CLAIMS

What is claimed is:

1. A biological chamber system comprising a biochamber and center lumen, said biochamber being defined by outer walls of an engineered Sertoli tissue construct.
2. The system according to claim 1, wherein said engineered Sertoli tissue construct form said center lumen surrounding a population of cells which are different than said engineered Sertoli tissue construct .
3. The system according to claim 2, wherein said center lumen contains pancreatic islet cells.
4. The system according to claim 2, wherein said center lumen contains neuronal cells.
5. The system according to claim 4, wherein said neuronal cells are NT2 neurons.
6. The system according to claim 1, wherein said outer walls are formed from a plurality of engineered Sertoli cells to form the tissue construct.
7. The system according to claim 6, wherein said outer walls comprise a immunoprotective system.
8. The system according to claim 1, wherein said biological chamber system is used for transplantation.

9. A transplantation facilitator comprising a biochamber.

10. The transplantation facilitator according to claim 9, wherein said biochamber is defined as having outer walls formed of engineered Sertoli tissue construct.

11. The transplantation facilitator according to claim 10, wherein said biochamber is further defined as having a center lumen surrounding therapeutic cells.

12. The transplantation facilitator according to claim 11, wherein said therapeutic cells are neuronal cells.

13. A method of making biochambers comprising the steps of:
co-culturing facilitator cells and therapeutic cells about the therapeutic cells to form a chamber thereabout; and
re-engineering the facilitator cells to form a tissue construct.

14. The method according to claim 13, further including the step of segregating the facilitator cells away from the therapeutic cells.

15. The method according to claim 14, wherein said segregating step further includes the step of inducing the epithelization and polarization of the facilitator cells.

16. The method according to claim 17, wherein said inducing step further includes adding a compound for inducing epithelization and polarization.

17. A method of transplanting cells comprising the steps of:
forming a biochamber,
incorporating therapeutic cells into said biochamber; and
transplanting the biochamber containing the transplant cells into a host

18. The method according to claim 17, wherein said incorporating step further includes co-culturing the therapeutic cells with cells which form the biochamber simultaneously.

19. A biochamber comprising an outer wall of facilitator cells and an inner lumen of therapeutic cells.

20. A biochamber comprising an outer wall of protective cells and an inner wall of secreting cells, said outer wall allowing for release from said biochamber of the secretions from said secreting cells.

21. A transplantation vessel comprising a housing made of one type of cell including an inner cavity and a center lumen surrounding a population of cells different from said housing.

22. The transplantation vessel according to claim 21, wherein said housing consists of engineered Sertoli tissue construct.


23. The transplantation vessel according to claim 21, wherein said cell population consists of at least one therapeutic cell.

24. The transplantation vessel according to claim 23, wherein said therapeutic cells are selected from the group consisting essentially of neuronal cells, NT2 cells, pancreatic islet cells, dopaminergic cells, and bovine chromaffin cells.

SERTOLI CELLS AS BIOCHAMBERS


ABSTRACT OF THE DISCLOSURE

According to the present invention, there is provided a biological chamber system having a biochamber defined by outer walls of Sertoli cells. Also provided is a transplantation facilitator including a biochamber. A method of making biochambers by co-culturing facilitator cells and therapeutic cells and then aggregating the facilitator cells is also provided. Also provided is a method of transplanting cells by incorporating transplant cells into a biochamber and transplanting the biochamber containing the transplant cells.



05/24/00 06:13:00


Conventional Culture



"Apical secretion in a closed compartment creates a fluid-filled lumen by appreciating hydrostatic pressure"

Blood-testis barrier"

Microgravity Coculture



Sertoli cells


Islets

(or NT₂ cells)




"Microgravity coculture results in the integration of therapeutic cells into Sertoli cell biochambers"

Fig 2




Positive FasL immunostaining identifies Sertoli cells and suggests a mechanism by which they may effect immune suppression at the graft site.



Activated T-lymphocytes with upregulated Fas receptors

FasL-Fas induced apoptosis of lymphocytes



The expression of FasL by Sertoli cells induces apoptosis of the invading immune cells by binding to the upregulated Fas receptors on these activated T-lymphocytes. This results in the attrition of these immune cells at the graft site thereby downregulating the immune responses-this by an already well-defined mechanism occurring naturally in the mammalian system.

09593629-061300

Isolated Sertoli cells from peripubertal rats and pancreatic islets from neonatal pigs were cocultured by conventional culture technology in the same medium described for the HARV simulated microgravity coculture. Sertoli cells were pre-plated 48 hours on plastic or Matrigel substrates. Pre-treated isolated pig islets were added to the Sertoli cell-enriched monoculture 24 hours later. This Sertoli-Islet coculture was incubated at 37°C and by 24 hr islets attached to and integrated into the underlying Sertoli cells. Within another 48-72hrs, Sertoli cells reorganized into spherical or chord-like aggregates. This process was enhanced for those cocultures in which Sertoli cells had been plated on the Matrigel. Islets appeared to retain their structural integrity better in the non-Matrigel cocultures (Fig 1) than in the cocultures not having a Matrigel substrate (Fig 2). Tissue constructs of Sertoli cells and pancreatic islet cells can be created in conventional coculture in a similar manner as that observed in simulated microgravity coculture.



Fig 1 Sertoli cell (SCs) and islets (arrows) in a Sertoli-islet tissue construct created in conventional coculture. B-cells are immunostained for insulin.

Fig 4





Fig 2. Sertoli cells (SCs) and B-cells (arrows) in a Sertoli-islet tissue construct created in conventional coculture, B-cells are immunostained for insulin.

Fig 5

1. Sanberg, P.R., C.V. Borlongan, A.I. Othberg, S. Saporta, T.B. Freeman and D.F. Cameron. Testis-derived Sertoli cells have a trophic effect on dopamine neurons and alleviate hemiparkinsonism in rats. *Mature medicine*, 3(10):1129-1132.

Figure 1.

1 Week HARV Coculture - rSCs + NT2




Sertoli-Neuron-Aggregate-Cells (SNACs) form *in vitro* following coculture of rat Sertoli cells and NT2 neuroes in simulated microgravity utilizing the High Aspect Rotation Velocity (HARV) bioreactor.

Fig 6


[illegible]

1 Week HARV Coculture - rSCs + hNT2 (neurons)

NT2 + SC
FasL



NT2 + SC
hNuMu



Immunocytochemical staining of mouse FasL and human nuclear matrix proteins in rSertoli-hNeuron Aggregated Cells (SNACs) following HARV incubated cocultures.

FIS 7

Docket No.
0152.00372

Declaration and Power of Attorney For Patent Application

English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

SERTOLI CELLS AS BIOCHAMBERS

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on _____ as United States Application No. or PCT International
Application Number _____
and was amended on _____

(if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

(Number)

(Country)

(Day/Month/Year Filed)

☐

(Number)

(Country)

(Day/Month/Year Filed)

☐

(Number)

(Country)

(Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

(Application Serial No.)

(Filing Date)

(Application Serial No.)

(Filing Date)

(Application Serial No.)

(Filing Date)

I hereby claim the benefit under 35 U. S. C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C. F. R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)
(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*

Kenneth I. Kohn, Reg. No. 30,955

Amy E. Rinaldo, Reg. No. 45,791

Send Correspondence to: **Amy E. Rinaldo**
KOHN & ASSOCIATES
30500 Northwestern Highway, Suite 410
Farmington Hills, Michigan 48334

Direct Telephone Calls to: *(name and telephone number)*

Amy E. Rinaldo (248) 539-5050

Full name of sole or first inventor

Don F. Cameron

Sole or first inventor's signature

Date

Residence

18206 Clear Lake Drive, Lutz, Florida 33549

Citizenship

U.S.

Post Office Address

Same as above.

Full name of second inventor, if any

Paul R. Sanberg

Second inventor's signature

Date

Residence

11751 Pilot Country Drive, Spring Hills, Florida 34610

Citizenship

U.S.

Post Office Address

Same as above.

Full name of third inventor, if any Samuel Saporta	
Third inventor's signature	Date
Residence 14026 Shady Shores Drive, Tampa, Florida 33613	
Citizenship U.S.	
Post Office Address Same as above.	

Full name of fourth inventor, if any Joel J. Hushen	
Fourth inventor's signature	Date
Residence 16914 Harrierridge Place, Lithia, Florida 33547	
Citizenship U.S.	
Post Office Address Same as above.	

Full name of fifth inventor, if any	
Fifth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	

Full name of sixth inventor, if any	
Sixth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	